

On first exit times for homogeneous diffusion processes*

Nikolai Dokuchaev[†]

In this paper we derive an upper bound for the expression $\mathbf{E}|T_1 - T_2|$, first exit times from a region for two homogeneous diffusion processes.

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a standard probability space and Q be a bounded simply connected region in $\mathbf{R}^n$. Consider the n - dimensional diffusion processes $y_i(t)$, $i=1,2$ (see [1, v.3, pp.257-258]) of the following Itô equations:

$$\begin{aligned} (1) \quad & dy_i(t) = f_i(y(t))dt + \beta_i(y(t))dw_t, \\ (2) \quad & y_i(0) = a_i. \end{aligned}$$

Here $t \geq 0$, w_t is a standard d -dimensional Wiener process, coordinated as usual with some right-continuous non-decreasing flow of σ -algebras $\mathcal{F}_t \subset \mathcal{F}$; f_i and β_i are nonrandom functions with respective values in $\mathbf{R}^n$ and $\mathbf{R}^{n \times d}$, (here and elsewhere $i = 1, 2$). The random vectors a_i are measurable with respect to the σ -algebra $\mathcal{F}_0$ and $a_i \in \bar{Q}$ with probability 1 ($\bar{Q}$ denotes the closure of the region Q). All vectors and matrices are real with Euclidean norm $|\cdot|$.

Consider the r.v.'s $T_i = \inf\{t : y_i(t) \notin Q\}$. These are the first exit times of the processes $y_i(t)$ from the region Q .

Consider in Q the Dirichlet problem

$$(3) \quad L_i v_i = -1, \quad v_i|_{\partial Q} = 0.$$

Here ∂Q is the boundary of Q and the differential operators

$$(4) \quad L_i = \sum_{j=1}^n f_i^{(j)} \frac{\partial}{\partial y^{(j)}} + \frac{1}{2} \sum_{j,k=1}^n b_i^{(j,k)} \frac{\partial^2}{\partial y^{(j)} \partial y^{(k)}},$$

where $f_i^{(j)}$, $y^{(j)}$, $b_i^{(j,k)}$ are the components of the vectors f_i , y and the matrices $b_i(\cdot) = \beta_i(\cdot)\beta_i(\cdot)^\top$.

We assume that all the components of the function f , β_i are continuously differentiable, the eigenvalues of the matrices b_i are isolated from zero uniformly in Q , and the boundary ∂Q is smooth. Problem (3)-(4) then has a (unique) solution that is twice continuously differentiable up to the boundary.

Denote by dv_i/dy the vector with components $\partial v_i / \partial y^{(j)}$.

*Theory of Probability and Its Application, 1987, v.31, No 3, pp. 497-498; English translation of the September 1986 issue of the Soviet journal "Teoria Veroyatnostei i ee primneniya"; translated from Russian by Amal Ellis

[†]Correspondence address: The Institute of Mathematics and Mechanics, St.Petersburg State University, Bibliotchnaya pl.2, Petrodvoretz, St.Petersburg, 198904, Russia. E-mail: dokuchaev@pobox.spbu.ru; fax 7(812)4286998

Theorem.

$$(5) \quad \mathbf{E}|T_1 - T_2| \leq \max_{i=1,2} \sup_{x \in Q} \left| \frac{dv_i}{dy}(y) \right| \mathbf{E}|y_1(\tilde{T}) - y_2(\tilde{T})|,$$

where $\tilde{T} = T_1 \wedge T_2 = \min(T_1, T_2)$.

Proof. Let e_1 and e_2 be the indicator functions of the events $\{T_1 > T_2\}$ and $\{T_2 > T_1\}$ respectively. Obviously,

$$(6) \quad \mathbf{E}|T_1 - T_2| = \mathbf{E}\{e_1(T_1 - T_2)\} + \mathbf{E}\{e_2(T_2 - T_1)\}.$$

The r.v.'s e_i are measurable with respect to the σ -algebras $\mathcal{F}_{\tilde{T}}$, $\mathcal{F}_{T_j}$, associated with the Markov times (with respect to the flow $\mathcal{F}_t$) $\tilde{T}$, T_j , $j = 1, 2$ (see [1, v.2, Chap.4, §2]). Using Itô's formula, we obtain the equality

$$(7) \quad \begin{aligned} \mathbf{E}\{e_1\{v_1[y_1(T_2)] - v_1[y_1(T_2)]\}\} &= -\mathbf{E}\{e_1\{v_1[y_1(T_1)] - v_1[y_1(T_2)]\}\} \\ &= -\mathbf{E}\left\{e_1 \int_{\tilde{T}}^{T_1} L_1 v_1[y_1(t)] dt\right\} = \mathbf{E}\{e_1(T_1 - T_2)\}. \end{aligned}$$

If we replaced the indices 1, 2 in (7) by 2, 1, we get an analogous expression for $\mathbf{E}\{e_1(T_1 - T_2)\}$. Now the assertion of the theorem follows at once from (6).

Example. In equations (1)-(2) let $n = d = 1$, $y_i(t) = a_i + w_t$, $Q = (0, 1)$. Then, obviously, $v_i(y) = y(1 - y)$, $dv_i(y)/dy = 1 - 2y$ and $\mathbf{E}|T_1 - T_2| \leq |a_1 - a_2|$.

The behavior of the first exit times of the process when the drift and diffusion coefficients vary has been studied in details in the book [2]. However, it solves problems of different sort than the one considered above.

For non-homogeneous processes and T_1, T_2 bounded times an estimate similar to (5) was derived in [3] and used to prove the "stochastic maximum principle" established in [4] for a controlled diffusion process in a bounded region. For strong Markov processes of arbitrary form, an estimate of type (5) was derived in [5] (however, the assumptions made there are restrictive and not easy to check for nondiffusional processes).

References

- [1] Gihman I.I. and Skorohod, A.V. *The Theory of Stochastic Processes*. Vol. 2, 1975; vol. 3, 1979. Springer-Verlag, New York, 1975.
- [2] A.D.Venttsel' and M.I.Freidlin, *Fluctuations in Dynamic Systems under the Action of Small Random Perturbations*, Nauka, Moscow, 1979 (in Russian).
- [3] Dokuchaev, N. On moment's of first emergency for diffusion-type processes. Leningrad Univ. Mechanical Bull. (1984), N 1, 58-61.
- [4] Dokuchaev, N. Optimal program control of a stochastic process cutting off on the boundary of a region. Working paper, Dep. VINITI No. 4705-83, 46 pp., 1983 (in Russian).
- [5] Dokuchaev, N. On the closeness of the first exit times for diffusion processes. Working paper, Dep. VINITI No. 5398-84, 8 pp., 1984 (in Russian).